

A short proof of Paouris' inequality

Radosław Adamczak ^{*} Rafał Łatała [†]

Alexander E. Litvak [‡] Krzysztof Oleszkiewicz [§]

Alain Pajor [¶] Nicole Tomczak-Jaegermann ^{||}

Abstract

We give a short proof of a result of G. Paouris on the tail behaviour of the Euclidean norm $|X|$ of an isotropic log-concave random vector $X \in \mathbb{R}^n$, stating that for every $t \geq 1$,

$$\mathbb{P}(|X| \geq ct\sqrt{n}) \leq \exp(-t\sqrt{n}).$$

More precisely we show that for any log-concave random vector X and any $p \geq 1$,

$$(\mathbb{E}|X|^p)^{1/p} \sim \mathbb{E}|X| + \sup_{z \in S^{n-1}} (\mathbb{E}|\langle z, X \rangle|^p)^{1/p}.$$

AMS Classification: 46B06, 46B09 (Primary), 52A23 (Secondary)

Key Words and Phrases: log-concave random vectors, deviation inequalities

^{*}Research partially supported by MNiSW Grant no. N N201 397437.

[†]Research partially supported by MNiSW Grant no. N N201 397437.

[‡]Research partially supported by the E.W.R. Steacie Memorial Fellowship.

[§]Research partially supported by MNiSW Grant no. N N201 397437.

[¶]Research partially supported by the ANR project ANR-08-BLAN-0311-01.

^{||}This author holds the Canada Research Chair in Geometric Analysis.

1 Introduction

Let X be a random vector in the Euclidean space \mathbb{R}^n equipped with its Euclidean norm $|\cdot|$ and its scalar product $\langle \cdot, \cdot \rangle$. Assume that X has a log-concave distribution (a typical example of such a distribution is a random vector uniformly distributed on a convex body). Assume further that it is centered and its covariance matrix is the identity – such a random vector will be called *isotropic*. A famous and important result of ([14], Theorem 1.1) states that

Theorem 1. *There exists an absolute constant $c > 0$ such that if X is an isotropic log-concave random vector in \mathbb{R}^n , then for every $t \geq 1$,*

$$\mathbb{P}(|X| \geq ct\sqrt{n}) \leq \exp(-t\sqrt{n}).$$

This result had a huge impact on the study of log-concave measures and has a lot of applications in that subject.

A Borel probability measure on \mathbb{R}^n is called log-concave if for all $0 < \theta < 1$ and all compact sets $A, B \subset \mathbb{R}^n$ one has

$$\mu((1 - \theta)A + \theta B) \geq \mu(A)^{1-\theta} \mu(B)^\theta.$$

We refer to [5, 6] for a general study of this class of measures. Clearly, the affine image of a log-concave probability is also log-concave. The Euclidean norm of an n -dimensional log-concave random vector has moments of all orders (see [5]). A log-concave probability is supported on some convex subset of an affine subspace where it has a density. In particular when the support of the probability generates the whole space \mathbb{R}^n (in which case we talk, in short, about full-dimensional probability) a characterization of Borell (see [5, 6]) states that the probability is absolutely continuous with respect to the Lebesgue measure and has a density which is log-concave. We say that a random vector is log-concave if its distribution is a log-concave measure.

Let $X \in \mathbb{R}^n$, be a random vector, denote the weak p -th moment of X by

$$\sigma_p(X) = \sup_{z \in S^{n-1}} (\mathbb{E}|\langle z, X \rangle|^p)^{1/p}.$$

The purpose of this article is to give a short proof of the following

Theorem 2. *For any log-concave random vector $X \in \mathbb{R}^n$ and any $p \geq 1$,*

$$(\mathbb{E}|X|^p)^{1/p} \leq C(\mathbb{E}|X| + \sigma_p(X)),$$

where C is an absolute positive constant.

This result may be deduced directly from Paouris' work in [14]. Indeed, it is a consequence of Theorem 8.2 combined with Lemma 3.9 in that paper. As formulated here, Theorem 2 first appeared in [3] (Theorem 2). Note that because trivially a converse inequality is valid (with constant 1/2), Theorem 2 states in fact an equivalence for $(\mathbb{E}|X|^p)^{1/p}$.

It is noteworthy that the following strengthening of Theorem 2 is still open: $(\mathbb{E}|X|^p)^{1/p} \leq \mathbb{E}|X| + C\sigma_p(X)$, where C is an absolute positive constant.

If X is a log-concave random vector, then so is $\langle z, X \rangle$ for every $z \in S^{n-1}$. It follows that there exists an absolute constant $C' > 0$ such that for any $p \geq 1$, $\sigma_p(X) \leq C'p\sigma_2(X)$ ([5]). (In fact one can deduce this inequality with $C' = 1$ from [4] or from Remark 5 in [11]; see also Remark 1 following Theorem 3.1 in [2].) If moreover X is isotropic, then $\mathbb{E}|X| \leq (\mathbb{E}|X|^2)^{1/2} = \sqrt{n}$ and $\sigma_2(X) = 1$; thus

$$(\mathbb{E}|X|^p)^{1/p} \leq C(\sqrt{n} + C'p).$$

From Markov's inequality for $p = t\sqrt{n}$, Theorem 2 implies Theorem 1 with $c = (C' + 1)eC$.

Let us recall the idea underlying the proof by Paouris. Let $X \in \mathbb{R}^n$ be an isotropic log-concave random vector. Let $p \sim \sqrt{n}$ be an integer (for example, $p = \lfloor \sqrt{n} \rfloor$). Let $Y = PX$ where P is an orthogonal projection of rank p and let $G \in \text{Im}P$ be a standard Gaussian vector. By rotation invariance, $\mathbb{E}|Y|^p \sim \mathbb{E}|\langle G/\sqrt{p}, Y \rangle|^p$. If the linear forms $\langle z, X \rangle$ with $|z| = 1$ had a sub-Gaussian tail behaviour, the proof would be straightforward. But in general they only obey a sub-exponential tail behaviour. The first step of the proof consists of showing that there exists some z for which $(\mathbb{E}|\langle z, Y \rangle|^p)^{1/p}$ is in fact small, compared to $\mathbb{E}|Y|$. The second step uses a concentration principle to show that $(\mathbb{E}_X|\langle z, PX \rangle|^p)^{1/p}$ is essentially constant on the sphere, for a random orthogonal projection of rank $p \sim \sqrt{n}$, and thus comparable to the minimum. Thus for these *good* projections, one has a good estimate of $(\mathbb{E}|Y|^p)^{1/p}$ and the result follows by averaging over P . Our proof follows the same scheme, at least for the first step, but whereas the proof of the first step in [14] is the most technical part, our argument is very simple. Then the estimate for $\min_{|z|=1} \mathbb{E}|\langle z, Y \rangle|^p$ brings us to a minimax problem precisely in the form answered by Gordon's inequality ([9]).

Finally we would like to note that our proof can be generalized to the case of convex measures in the sense of [5, 6]. Of course the proof is longer and more technical. We provide the details in [1].

2 Proof of Theorem 2

First let us notice that it is enough to prove Theorem 2 for symmetric log-concave random vectors. Indeed, let X be a log-concave random vector and let X' be an independent copy. By Jensen's inequality we have for all $p \geq 1$,

$$(\mathbb{E}|X|^p)^{1/p} \leq (\mathbb{E}|X - \mathbb{E}X|^p)^{1/p} + |\mathbb{E}X| \leq (\mathbb{E}|X - X'|^p)^{1/p} + \mathbb{E}|X|.$$

On the other hand $\mathbb{E}|X - X'| \leq 2\mathbb{E}|X|$ and for $p \geq 1$ one has $\sigma_p(X - X') \leq 2\sigma_p(X)$. Since $X - X'$ is log-concave (see [8]) and symmetric, we obtain that the symmetric case proved with a constant C' implies the non-symmetric case with the constant $C = 2C' + 1$.

Lemma 3. Let $Y \in \mathbb{R}^q$ be a random vector. Let $\|\cdot\|$ be a norm on \mathbb{R}^q . Then for all $p > 0$,

$$\min_{|z|=1} (\mathbb{E}|\langle z, Y \rangle|^p)^{1/p} \leq \frac{(\mathbb{E}\|Y\|^p)^{1/p}}{\mathbb{E}\|Y\|} \mathbb{E}|Y|.$$

Proof: Let r be the largest number such that $r\|t\| \leq |t|$ for all $t \in \mathbb{R}^q$. Using duality, pick $z \in \mathbb{R}^q$ such that $|z| = 1$ and $\|z\|_* \leq r$ (the dual norm of $\|\cdot\|$). Then $|\langle z, t \rangle| \leq r\|t\| \leq |t|$ for all $t \in \mathbb{R}^q$. Therefore, $(\mathbb{E}|\langle z, Y \rangle|^p)^{1/p} \leq r(\mathbb{E}\|Y\|^p)^{1/p}$ for any $p > 0$, and the proof follows from $r\mathbb{E}\|Y\| \leq \mathbb{E}|Y|$. \square

Lemma 4. Let Y be a full-dimensional symmetric log-concave \mathbb{R}^q -valued random vector. Then there exists a norm $\|\cdot\|$ on \mathbb{R}^q such that

$$(\mathbb{E}\|Y\|^q)^{1/q} \leq 500 \mathbb{E}\|Y\|.$$

Remark. In fact the constant 500 can be significantly improved. To keep the presentation short and transparent we omit the details.

Proof: From Borell's characterization Y has an even log-concave density g_Y . Thus $g_Y(0)$ is the maximum of g_Y . Define a symmetric convex set by

$$K = \{t \in \mathbb{R}^q : g_Y(t) \geq 25^{-q} g_Y(0)\}.$$

Since clearly K has a non-empty interior, it is the unit ball of a norm which we denote by $\|\cdot\|$. On one hand, $1 \geq \mathbb{P}(Y \in K) = \int_K g_Y \geq 25^{-q} g_Y(0) \text{vol}(K)$, thus

$$\mathbb{P}(\|Y\| \leq 1/50) = \int_{K/50} g_Y \leq g_Y(0) 50^{-q} \text{vol}(K) \leq 2^{-q} \leq 1/2.$$

Therefore $\mathbb{E}\|Y\| \geq \mathbb{P}(\|Y\| > 1/50)/50 \geq 1/100$. On the other hand, by the log-concavity of g_Y ,

$$\forall t \in \mathbb{R}^q \setminus K \quad g_{2Y}(t) = 2^{-q} g_Y(t/2) \geq 2^{-q} g_Y(t)^{1/2} g_Y(0)^{1/2} \geq (5/2)^q g_Y(t).$$

Therefore

$$\mathbb{E}\|Y\|^q \leq 1 + \mathbb{E}(\|Y\|^q 1_{Y \in \mathbb{R}^q \setminus K}) \leq 1 + (2/5)^q \mathbb{E}\|2Y\|^q = 1 + (4/5)^q \mathbb{E}\|Y\|^q.$$

We conclude that $(\mathbb{E}\|Y\|^q)^{1/q} \leq 5$ and $(\mathbb{E}\|Y\|^q)^{1/q} / \mathbb{E}\|Y\| \leq 500$. \square

Lemma 5. Let $n, q \geq 1$ be integers and $p \geq 1$. Let X be an n -dimensional random vector, G be a standard Gaussian vector in \mathbb{R}^n and Γ be an $n \times q$ standard Gaussian matrix. Then

$$(\mathbb{E}|X|^p)^{1/p} \leq \alpha_p^{-1} \left(\mathbb{E} \min_{|t|=1} \|\Gamma t\| + (\alpha_p + \sqrt{q}) \sigma_p(X) \right),$$

where $\|z\| = (\mathbb{E}|\langle z, X \rangle|^p)^{1/p}$ and α_p^p is the p -th moment of an $N(0, 1)$ Gaussian random variable (so that $\lim_{p \rightarrow \infty} (\alpha_p / \sqrt{p}) = 1/\sqrt{e}$).

Proof: By rotation invariance, $\mathbb{E}|\langle G, X \rangle|^p = \alpha_p^p \mathbb{E}|X|^p$. Notice that

$$\sigma^2 := \sup_{|||t|||_* \leq 1} \mathbb{E}|\langle G, t \rangle|^2 = \sup_{|||t|||_* \leq 1} |t|^2 = \sigma_p^2(X),$$

where $|||\cdot|||_*$ denotes the norm on \mathbb{R}^n dual to the norm $|||\cdot|||$. Denote the median of $|||G|||$ by M_G . The classical deviation inequality for a norm of a Gaussian vector ([7], [15], see also [12], Theorem 12.2) states

$$\forall s \geq 0 \quad \mathbb{P}(|||G||| - M_G| \geq s) \leq 2 \int_{s/\sigma}^{\infty} \exp(-t^2/2) \frac{dt}{\sqrt{2\pi}}$$

and since $M_G \leq \mathbb{E}|||G|||$ ([10], see also [12], Lemma 12.2) this implies

$$(\mathbb{E}|X|^p)^{1/p} = \alpha_p^{-1} (\mathbb{E}|||G|||^p)^{1/p} \leq \alpha_p^{-1} (\mathbb{E}|||G||| + \alpha_p \sigma_p(X))$$

(cf. [13], Statement 3.1).

The Gordon minimax lower bound (see [9], Theorem 2.5) states that for any norm $|||\cdot|||$

$$\mathbb{E}|||G||| \leq \mathbb{E} \min_{|t|=1} |||\Gamma t||| + (\mathbb{E}|H|) \max_{|z|=1} |||z||| \leq \mathbb{E} \min_{|t|=1} |||\Gamma t||| + \sqrt{q} \sigma_p(X),$$

where H is a standard Gaussian vector in \mathbb{R}^q . This concludes the proof. \square

Proof of Theorem 2: Assume that X is log-concave symmetric. We use the notation of Lemma 5 with q the integer such that $p \leq q < p+1$. We first condition on Γ . Let $Y = \Gamma^* X$. Note that Y is log-concave symmetric and that

$$|||\Gamma t||| = (\mathbb{E}_X |\langle \Gamma t, X \rangle|^p)^{1/p} = (\mathbb{E}_X |\langle t, \Gamma^* X \rangle|^p)^{1/p}.$$

If $\Gamma^* X$ is supported by a hyperplane then $\min_{|t|=1} (\mathbb{E}_X |\langle t, \Gamma^* X \rangle|^p)^{1/p} = 0$. Otherwise Lemma 4 applies and combined with Lemma 3 gives that

$$\min_{|t|=1} |||\Gamma t||| \leq \min_{|t|=1} (\mathbb{E}_X |\langle t, \Gamma^* X \rangle|^p)^{1/p} \leq 500 \mathbb{E}_X |\Gamma^* X|.$$

By taking expectation over Γ we get

$$\mathbb{E} \min_{|t|=1} |||\Gamma t||| \leq 500 \mathbb{E} |\Gamma^* X| = 500 \mathbb{E} |H| \mathbb{E} |X| \leq 500 \sqrt{q} \mathbb{E} |X|,$$

where $H \in \mathbb{R}^q$ is a standard Gaussian vector. Applying Lemma 5 we obtain

$$(\mathbb{E}|X|^p)^{1/p} \leq 500 \alpha_p^{-1} \sqrt{q} \mathbb{E}|X| + (1 + \alpha_p^{-1} \sqrt{q}) \sigma_p(X).$$

This implies the desired result, since $q \leq p+1$ and hence $\alpha_p^{-1} \sqrt{q} \leq c$ for some numerical constant c (recall that $\lim_{p \rightarrow \infty} (\alpha_p / \sqrt{p}) = 1/\sqrt{e}$). \square

References

- [1] R. Adamczak, O. Guédon, R. Latała, A. E. Litvak, K. Oleszkiewicz, A. Pajor and N. Tomczak-Jaegermann, *Moment estimates for convex measures*, preprint.
- [2] R. Adamczak, R. Latała, A. E. Litvak, A. Pajor and N. Tomczak-Jaegermann, *Tail estimates for norms of sums of log-concave random vectors*, preprint, <http://arxiv.org/abs/1107.4070>.
- [3] R. Adamczak, R. Latała, A. E. Litvak, A. Pajor and N. Tomczak-Jaegermann, *Geometry of log-concave Ensembles of random matrices and approximate reconstruction*, C.R. Math. Acad. Sci. Paris, **349** (2011), 783–786.
- [4] R. E. Barlow, A. W. Marshall and F. Proschan, *Properties of probability distributions with monotone hazard rate*, Ann. Math. Statist., **34** (1963), 375–389.
- [5] C. Borell, *Convex measures on locally convex spaces*, Ark. Math., **12** (1974), 239–252.
- [6] C. Borell, *Convex set functions in d -space*, Period. Math. Hungar., **6** (1975), 111–136.
- [7] C. Borell, *The Brunn-Minkowski inequality in Gauss space*, Invent. Math., **30** (1975), 207–216.
- [8] Ju. S. Davidovic, B. I. Korenbljum and B. I. Hacet, *A certain property of logarithmically concave functions*, Soviet Math. Dokl., **10** (1969), 447–480; translation from Dokl. Akad. Nauk SSSR **185** (1969), 1215–1218.
- [9] Y. Gordon, *Some inequalities for Gaussian processes and applications*, Israel J. Math., **50** (1985), 265–289.
- [10] S. Kwapień, *A remark on the median and the expectation of convex functions of Gaussian vectors*, Probability in Banach spaces, 9 (Sandjberg, 1993), 271–272, Progr. Probab., 35, Birkhuser Boston, Boston, MA, 1994.
- [11] S. Kwapień, R. Latała and K. Oleszkiewicz, *Comparison of moments of sums of independent random variables and differential inequalities*, J. Funct. Anal., **136** (1996), 258–268.
- [12] M. A. Lifshits, *Gaussian random functions*. Mathematics and its Applications; 322. Kluwer Academic Publishers, Dordrecht, 1995.
- [13] A. E. Litvak, V. D. Milman and G. Schechtman, *Averages of norms and quasi-norms*, Math. Ann., **312** (1998), 95–124.
- [14] G. Paouris, *Concentration of mass on convex bodies*, Geom. Funct. Anal., **16** (2006), 1021–1049.

- [15] V. N. Sudakov, B. S. Cirel'son, *Extremal properties of half-spaces for spherically invariant measures*, J. Sov. Math. **9** (1978), 9–18; translation from Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov (LOMI) **41** (1974), 14–24.

Radosław Adamczak,
Institute of Mathematics,
University of Warsaw,
Banacha 2, 02-097 Warszawa, Poland
e-mail: `radamcz@mimuw.edu.pl`

Rafał Łatała,
Institute of Mathematics,
University of Warsaw,
Banacha 2, 02-097 Warszawa, Poland
e-mail: `rlatala@mimuw.edu.pl`

Alexander E. Litvak,
Dept. of Math. and Stat. Sciences,
University of Alberta,
Edmonton, Alberta, Canada, T6G 2G1.
e-mail: `alexandr@math.ualberta.ca`

Krzysztof Oleszkiewicz,
Institute of Mathematics,
University of Warsaw,
Banacha 2, 02-097 Warszawa, Poland
e-mail: `koles@mimuw.edu.pl`

Alain Pajor,
Université Paris-Est
Équipe d'Analyse et Mathématiques Appliquées,
5, boulevard Descartes, Champs sur Marne,
77454 Marne-la-Vallée, Cedex 2, France
e-mail: `Alain.Pajor@univ-mlv.fr`

Nicole Tomczak-Jaegermann,
Dept. of Math. and Stat. Sciences,
University of Alberta,
Edmonton, Alberta, Canada, T6G 2G1.
e-mail: `nicole.tomczak@ualberta.ca`